

## Bijlage 16. Referenties

- Abu-Mostafa NA, Alqahtani A, Abu-Hasna M, et al: A randomized clinical trial compared the effect of intra-alveolar 0.2% chlorhexidine bio-adhesive gel versus 0.12% chlorhexidine rinse in reducing alveolar osteitis following molar teeth extractions. *Med Oral Patol Oral Cir Bucal* 20:e82, 2015
- Agbaje JO, Heijsters G, Salem AS, Van Slycke S, Schepers S, Politis C, Vrielinck L. Coronectomy of Deeply Impacted Lower Third Molar: Incidence of Outcomes and Complications after One Year Follow-Up. *J Oral Maxillofac Res.* 2015 Jun 30;6(2):e1.
- Al-Asfour A. Postoperative infection after surgical removal of impacted mandibular third molars: an analysis of 110 consecutive procedures. *Med Princ Pract.* 2009;18(1):48-52.
- Allen RT, Witherow H, Collyer J, Roper-Hall R, Nazir MA, Mathew G. The mesioangular third molar--to extract or not to extract? Analysis of 776 consecutive third molars. *Br Dent J.* 2009 Jun 13;206(11):E23
- Andrews J (1), Guyatt G, Oxman AD, Alderson P, Dahm P, Falck-Ytter Y, Nasser M, Meerpohl J, Post PN, Kunz R, Brozek J, Vist G, Rind D, Akl EA, Schünemann HJ. GRADE guidelines: 14. Going from evidence to recommendations: the significance and presentation of recommendations. *J Clin Epidemiol.* 2013 Jul;66(7):719-25.
- Andrews JC (2), Schünemann HJ, Oxman AD, Pottie K, Meerpohl JJ, Coello PA, Rind D, Montori VM, Brito JP, Norris S, Elbarbary M, Post P, Nasser M, Shukla V, Jaeschke R, Brozek J, Djulbegovic B, Guyatt G. GRADE guidelines: 15. Going from evidence to recommendation--determinants of a recommendation's direction and strength. *J Clin Epidemiol.* 2013 Jul;66(7):726-35.
- Anjrini AA, Kruger E, Tennant M. Cost effectiveness modelling of a 'watchful monitoring strategy' for impacted third molars vs prophylactic removal under GA: an Australian perspective. *Br Dent J.* 2015 Jul 10;219(1):19-23.
- Arteagoitia I, Diez A, Barbier L, Santamaria G, Santamaria J. Efficacy of amoxicillin/clavulanic acid in preventing infectious and inflammatory complications following impacted mandibular third molar extraction. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology* 2005;100(1):e11-8.
- Babar A, Ibrahim MW, Baig NJ, et al: Efficacy of intra-alveolar chlorhexidine gel in reducing frequency of alveolar osteitis in mandibular third molar surgery. *J Coll Physicians Surg Pak* 22: 91, 2012
- Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, Vist GE, Falck-Ytter Y, Meerpohl J, Norris S, Guyatt GH. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol.* 2011 Apr;64(4):401-6.
- Baqain ZH, Al-Shafii A, Hamdan AA, Sawair FA. Flap design and mandibular third molar surgery: A split mouth randomized clinical study. *International Journal of Oral and Maxillofacial Surgery* 2012; Vol. 41, issue 8:1020-4.
- Baqain ZH, Karaky AA, Sawair F, Khraisat A, Duaibis R, Rajab LD. Frequency estimates and risk factors for postoperative morbidity after third molar removal: a prospective cohort study. *J Oral Maxillofac Surg.* 2008 Nov;66(11):2276-83.
- Barclay JK. Metronidazole and dry socket: prophylactic use in mandibular third molar removal complicated by nonacute pericoronitis. *New Zealand Dental Journal* 1987;83: 71-5.
- Barone A, Marconcini S, Giacomelli L, Rispoli L, Calvo JL, Covani U. A randomized clinical evaluation of ultrasound bone surgery versus traditional rotary instruments in lower third molar extraction. *Journal of Oral and Maxillofacial Surgery* 2010; Vol. 68, issue 2:330-6.
- Bastos Carmo A do, de Oliveira JB, Mello KF, Leão PB, Artese F, Normando D. The ability of orthodontists and oral/maxillofacial surgeons to predict eruption of lower third molar. *Prog Orthod.* 2016 Dec;17(1):21.
- Bello SA, Olaitan AA, Ladeinde AL. A randomized comparison of the effect of partial and total wound closure techniques on postoperative morbidity after mandibular third molar surgery. *Journal of Oral and Maxillofacial Surgery* 2011;69:e24-30.
- Benediktsdóttir IS, Wenzel A, Petersen JK, Hintze H. Mandibular third molar removal: risk indicators for extended operation time, postoperative pain, and complications. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004 Apr;97(4):438-46.
- Bergdahl M, Hedstrom L. Metronidazole for the prevention of dry socket after removal of partially impacted mandibular third molar: a randomised controlled trial. *British Journal of Oral & Maxillofacial Surgery* 2004;42(6):555-8.
- Berwick JE, Lessin ME: Effects of a chlorhexidine gluconate oral rinse on the incidence of alveolar osteitis in mandibular third molar surgery. *J Oral Maxillofac Surg* 48:444, 1990
- Bienstock DA, Dodson TB, Perrott DH, Chuang SK. Prognostic factors affecting the duration of disability after third molar removal. *J Oral Maxillofac Surg.* 2011 May;69(5):1272-7.
- Blakey GH, Marciani RD, Haug RH, Phillips C, Offenbacher S, Pabla T, White RP Jr. Periodontal pathology associated with asymptomatic third molars. *J Oral Maxillofac Surg.* 2002 Nov;60(11):1227-33.
- Blondeau F, Daniel NG. Extraction of impacted mandibular third molars: postoperative complications and their risk factors. *J Can Dent Assoc.* 2007 May;73(4):325.
- Bortoluzzi MC, Capella DL, Barbieri T, et al: A single dose of amoxicillin and dexamethasone for prevention of postoperative complications in third molar surgery: A randomized, doubleblind, placebo controlled clinical trial. *J Clin Med Res* 5:26, 2013.
- Briguglio F, Zenobio EG, Isola G, Briguglio R, Briguglio E, Farronato D, et al. Complications in surgical removal of impacted mandibular third molars in relation to flap design: clinical and statistical evaluations. *Quintessence International* 2011; Vol. 42, issue 6:445-53.
- Butler DP, Sweet JB. Effect of lavage on the incidence of localized osteitis in mandibular third molar extraction sites. *Oral Surgery* 1977;44(1):14-20.
- Bystedt H, von Konow L, Nord CE. A comparison of the effect of phenoxymethylpenicillin and azidocillin on postoperative complications after surgical removal of impacted mandibular third molars. *Swedish Dental Journal* 1981;5(5-6):225-34.
- Carter K, Worthington S. Predictors of third molar impaction: a systematic review and meta-analysis. *J Dent Res* 2016; 95: 267-76.
- Cerqueira PR, Vasconcelos BC, Bessa-Nogueira RV. Comparative study of the effect of a tube drain in impacted lower third molar surgery. *Journal of Oral and Maxillofacial Surgery* 2004;62(1):57-61.

- Chang SW, Shin SY, Kum KY, Hong J. Correlation study between distal caries in the mandibular second molar and the eruption status of the mandibular third molar in the Korean population. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2009 Dec;108(6):838-43.
- Cheung LK, Leung YY, Chow LK, Wong MC, Chan EK, Fok YH. Incidence of neurosensory deficits and recovery after lower third molar surgery: a prospective clinical study of 4338 cases. *Int J Oral Maxillofac Surg*. 2010 Apr;39(4):320-6.
- Chu FC, Li TK, Lui VK, Newsome PR, Chow RL, Cheung LK. Prevalence of impacted teeth and associated pathologies--a radiographic study of the Hong Kong Chinese population. *Hong Kong Med J*. 2003 Jun;9(3):158-63.
- Chuang SK, Perrott DH, Susarla SM, Dodson TB. Age as a risk factor for third molar surgery complications. *J Oral Maxillofac Surg*. 2007 Sep;65(9):1685-92.
- Chukwunke FN, Oji C, Saheeb DB. A comparative study of the effect of using a rubber drain on postoperative discomfort following lower third molar surgery. *International Journal of Oral and Maxillofacial Surgery* 2008; 37(4):341-4.
- Clé-Ovejero A, Sánchez-Torres A, Camps-Font O, Gay-Escoda C, Figueiredo R, Valmaseda-Castellón E. Does 3-dimensional imaging of the third molar reduce the risk of experiencing inferior alveolar nerve injury owing to extraction?: A meta-analysis. *J Am Dent Assoc*. 2017 Aug;148(8):575-583.
- Danda AK, Krishna Tatiparthi M, Narayanan V, Siddareddi A. Influence of primary and secondary closure of surgical wound after impacted mandibular third molar removal on postoperative pain and swelling--a comparative and split mouth study. *Journal of Oral and Maxillofacial Surgery* 2010;68(2):309-12.
- de Brabander EC, Cattaneo G. The effect of surgical drain together with a secondary closure technique on postoperative trismus, swelling and pain after mandibular third molar surgery. *International Journal of Oral and Maxillofacial Surgery* 1998;17(2):119-21.
- Delilbasi C, Saracoglu SH, Kesken A: Effects of 0.2% chlorhexidine gluconate and amoxicillin plus clavulanic acid on the prevention of alveolar osteitis following mandibular third molar extractions. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 94:301, 2002
- Divaris K, Fisher EL, Shugars DA, White RP Jr. Risk factors for third molar occlusal caries: a longitudinal clinical investigation. *J Oral Maxillofac Surg*. 2012 Aug;70(8):1771-80.
- Doğan N, Orhan K, Günaydin Y, Köymen R, Okçu K, Uçok O. Unerupted mandibular third molars: symptoms, associated pathologies, and indications for removal in a Turkish population. *Quintessence Int*. 2007 Sep;38(8):e497-505.
- Dolanmaz D, Yildirim G, Isik K, Kucuk K, Ozturk A. A preferable technique for protecting the inferior alveolar nerve: coronectomy. *J Oral Maxillofac Surg*. 2009;67:1234-8.
- Elter JR, Offenbacher S, White RP, Beck JD. Third molars associated with periodontal pathology in older Americans. *J Oral Maxillofac Surg*. 2005 Feb;63(2):179-84.
- Erdogan O, Tatli U, Ustun Y, Damlar I. Influence of two different flap designs on the sequelae of mandibular third molar surgery. *Oral and Maxillofacial Surgery* 2011;15(3): 147-52.
- Falci SG, de Castro CR, Santos RC, de Souza Lima LD, Ramos-Jorge ML, Botelho AM, Dos Santos CR. Association between the presence of a partially erupted mandibular third molar and the existence of caries in the distal of the second molars. *Int J Oral Maxillofac Surg*. 2012 Oct;41(10):1270-4.
- Fisher EL, Garaas R, Blakey GH, Offenbacher S, Shugars DA, Phillips C, White RP Jr. Changes over time in the prevalence of caries experience or periodontal pathology on third molars in young adults. *J Oral Maxillofac Surg*. 2012 May;70(5):1016-22.
- Freudenthal N, Sternudd M, Jansson L, et al: A double-blind randomized study evaluating the effect of intra-alveolar chlorhexidine gel on alveolar osteitis after removal of mandibular third molars. *J Oral Maxillofac Surg* 73:600, 2015
- Garaas RN, Fisher EL, Wilson GH, Phillips C, Shugars DA, Blakey GH, Marciani RD, White RP Jr. Prevalence of third molars with caries experience or periodontal pathology in young adults. *J Oral Maxillofac Surg*. 2012 Mar;70(3):507-13.
- Gargallo-Albiol J, Buenechea-Imaz R, Gay-Escoda C. Lingual nerve protection during surgical removal of lower third molars. A prospective randomised study. *International Journal of Oral and Maxillofacial Surgery* 2000;29(4):268-71.
- Ghaemina H, Gerlach NL, Hoppenreijts TJ, Kicken M, Dings JP, Borstlap WA, de Haan T, Bergé SJ, Meijer GJ, Maal TJ. Clinical relevance of cone beam computed tomography in mandibular third molar removal: A multicentre, randomised, controlled trial. *J Craniomaxillofac Surg*. 2015 Dec;43(10):2158-67.
- Ghaemina H, Hoppenreijts TJ, Xi T, Fennis JP, Maal TJ, Bergé SJ, Meijer GJ. Postoperative socket irrigation with drinking tap water reduces the risk of inflammatory complications following surgical removal of third molars: a multicenter randomized trial. *Clin Oral Investig*. 2017 Jan;21(1):71-83
- Ghaemina H, Meijer GJ, Soehardi A, Borstlap WA, Mulder J, Bergé SJ. Position of the impacted third molar in relation to the mandibular canal. Diagnostic accuracy of cone beam computed tomography compared with panoramic radiography. *Int J Oral Maxillofac Surg*. 2009 Sep;38(9):964-71.
- Ghaemina H, Perry J, Nienhuijs MEL, Toedtling V, Tummers M, Hoppenreijts TJM, Van der Sanden WJM, Mettes TG. Surgical removal versus retention for the management of asymptomatic disease-free impacted wisdom teeth. *Cochrane Database of Systematic Reviews* 2016, Issue 8. Art. No.: CD003879.
- Goldsmith SM, De Silva RK, Tong DC, Love RM. Influence of a pedicle flap design on acute postoperative sequelae after lower third molar removal. *International Journal of Oral and Maxillofacial Surgery* 2012;41(3):371-5.
- Gomes ACA, Vasconcelos BC, de Oliveira e Silva ED, da Silva LC. Lingual nerve damage after mandibular third molar surgery: a randomised clinical trial. *Journal of Oral and Maxillofacial Surgery* 2005;63(10):1443-6.
- Goto S, Kurita K, Kuroiwa Y, Hatano Y, Kohara K, Izumi M, et al. Clinical and dental computed tomographic evaluation 1 year after coronectomy. *J Oral Maxillofac Surg*. 2012;70:1023-9.
- Greenwood M, Langton SG, Rood JP. A comparison of broad and narrow retractors for lingual nerve protection during lower third molar surgery. *British Journal of Oral and Maxillofacial Surgery* 1994;32(2):114-7.

- Grossi GB, Maiorana C, Garramone RA, Borgonovo A, Creminelli L, Santoro F. Assessing postoperative discomfort after third molar surgery: a prospective study. *J Oral Maxillofac Surg.* 2007 May;65(5):901-17.
- Guerrero ME, Botetano R, Beltran J, Horner K, Jacobs R. Can preoperative imaging help to predict postoperative outcome after wisdom tooth removal? A randomized controlled trial using panoramic radiography versus cone-beam CT. *Clin Oral Investig.* 2014 Jan;18(1):335-42.
- Güven O, Keskin A, Akal UK. The incidence of cysts and tumors around impacted third molars. *Int J Oral Maxillofac Surg.* 2000 Apr;29(2):131-5.
- Guyatt G (1), Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, Norris S, Falck-Ytter Y, Glasziou P, De Beer H, Jaeschke R, Rind D, Meerpohl J, Dahm P, Schünemann HJ. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol.* 2011 Apr;64(4):383-94.
- Guyatt GH (2), Oxman AD, Kunz R, Atkins D, Brozek J, Vist G, Alderson P, Glasziou P, Falck-Ytter Y, Schünemann HJ. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol.* 2011 Apr;64(4):395-400.
- Halpern LR, Dodson TB. Does prophylactic administration of systemic antibiotics prevent postoperative inflammatory complications after third molar surgery?. *Journal of Oral and Maxillofacial Surgery* 2007;65(2):177-85.
- Hanna K, Sambrook P, Armfield JM, Brennan DS. Preferences for dental decisional control and associations with quality of life among third molar patients attending public dental services. *Community Dent Health.* 2017 Sep;34(3):163-168.
- Happonen RP, Backstrom AC, Ylipaavalniemi P. Prophylactic use of phenoxymethylpenicillin and tinidazole in mandibular third molar surgery, a comparative placebo controlled clinical trial. *British Journal of Oral & Maxillofacial Surgery* 1990;28(1):12-5.
- Harada N, Beloor Vasudeva S, Matsuda Y, Seki K, Kapila R, Ishikawa N, Okano T, Sano T. Characteristic findings on panoramic radiography and cone-beam CT to predict paresthesia after extraction of impacted third molar. *Bull Tokyo Dent Coll.* 2015;56(1):1-8.
- Haraji A, Motamedi MH, Rezvani F. Can flap design influence the incidence of alveolar osteitis following removal of impacted mandibular third molars?. *General Dentistry* 2010; Vol. 58, issue 5:e187-9.
- Haraji A, Rakhshan V, Khamverdi N, et al: Effects of intra-alveolar placement of 0.2% chlorhexidine bioadhesive gel on dry socket incidence and postsurgical pain: A double-blind split-mouth randomized controlled clinical trial. *J Orofac Pain* 27:256, 2013
- Harradine NW, Pearson MH, Toth B. The effect of extraction of third molars on late lower incisor crowding: a randomized controlled trial. *British Journal of Orthodontics* 1998;25(2):117-22.
- Hasani A, Ahmadi Moshtaghin F, Roohi P, Rakhshan V. Diagnostic value of cone beam computed tomography and panoramic radiography in predicting mandibular nerve exposure during third molar surgery. *Int J Oral Maxillofac Surg.* 2016 Oct 31. pii: S0901-5027(16)30276-4.
- Hasegawa T, Ri S, Shigeta T, Akashi M, Imai Y, Kakei Y, Shibuya Y, Komori T. Risk factors associated with inferior alveolar nerve injury after extraction of the mandibular third molar--a comparative study of preoperative images by panoramic radiography and computed tomography. *Int J Oral Maxillofac Surg.* 2013 Jul;42(7):843-51.
- Hasegawa T, Ri S, Umeda M, Komori T. Multivariate relationships among risk factors and hypoesthesia of the lower lip after extraction of the mandibular third molar. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2011 Jun;111(6):e1-7.
- Hasegawa Y, Terada K, Kageyama I, Tsuchimochi T, Ishikawa F, Nakahara S. Influence of third molar space on angulation and dental arch crowding. *Odontology.* 2013 Jan;101(1):22-8.
- Hashemi HM, Beshkar M, Aghajani R. The effect of sutureless wound closure on postoperative pain and swelling after impacted mandibular third molar surgery. *British Journal of Oral and Maxillofacial Surgery* 2012;50(3):256-8.
- Haug RH, Perrott DH, Gonzalez ML, Talwar RM. The American Association of Oral and Maxillofacial Surgeons Age-Related Third Molar Study. *J Oral Maxillofac Surg.* 2005 Aug;63(8):1106-14.
- Hazza'a AM(1), Bataineh AB, Odat AA. Angulation of mandibular third molars as a predictive factor for pericoronitis. *J Contemp Dent Pract.* 2009 May 1;10(3):51-8.
- Hermesch CB, Hilton TJ, Biesbrock AR, et al: Perioperative use of 0.12% chlorhexidine gluconate for the prevention of alveolar osteitis: Efficacy and risk factor analysis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 85:381, 1998
- Hita-Iglesias P, Torres-Lagares D, Flores-Ruiz R, et al: Effectiveness of chlorhexidine gel versus chlorhexidine rinse in reducing alveolar osteitis in mandibular third molar surgery. *J Oral Maxillofac Surg* 66:441, 2008
- Huang GJ, Cunha-Cruz J, Rothen M, Spiekerman C, Drangsholt M, Anderson L, Roset GA. A prospective study of clinical outcomes related to third molar removal or retention. *Am J Public Health.* 2014 Apr;104(4):728-34.
- Kaczmarzyk T, Wichlinski J, Stypulkowska J, Zaleska M, Panas M, Woron J. Single-dose and multi-dose clindamycin therapy fails to demonstrate efficacy in preventing infectious and inflammatory complications in third molar surgery. *International Journal of Oral and Maxillofacial Surgery* 2007; 36(5):417-22.
- Kang F, Huang C, Sah MK, Jiang B. Effect of Eruption Status of the Mandibular Third Molar on Distal Caries in the Adjacent Second Molar. *J Oral Maxillofac Surg.* 2016 Apr;74(4):684-92.
- Kirk DG, Liston PN, Tong DC, Love RM. Influence of two different flap designs on incidence of pain, swelling, trismus, and alveolar osteitis in the week following third molar surgery. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* 2007;104(1):e1-6.
- Kouwenberg AJ, Stroy LP, Rijdt ED, Mensink G, Gooris PJ. Coronectomy of the mandibular third molar: Respect for the inferior alveolar nerve. *J Craniomaxillofac Surg.* 2016 May;44(5):616-21.
- Krekmanov L, Hallander HO. Relationship between bacterial contamination and alveolitis after third molar surgery. *International Journal of Oral Surgery* 1980;9(4): 274-80.
- Krekmanov L, Nordenram A: Postoperative complications after surgical removal of mandibular third molars. Effects of penicillin V and chlorhexidine. *Int J Oral Maxillofac Surg* 15:25, 1986

- Krekmanov L. Alveolitis after operative removal of third molars in the mandible. *International Journal of Oral Surgery* 1981;10(3):173-9.
- Kugelberg CF, Ahlström U, Ericson S, Hugoson A, Thilander H. The influence of anatomical, pathophysiological and other factors on periodontal healing after impacted lower third molar surgery. A multiple regression analysis. *J Clin Periodontol.* 1991 Jan;18(1):37-43.
- Lacasa JM, Jiménez JA, Ferrás V, Bossom M, Sóla-Morales O, García-Rey C, et al. Prophylaxis versus pre-emptive treatment for infective and inflammatory complications of surgical third molar removal: a randomized, double-blind, placebo-controlled, clinical trial with sustained release amoxicillin/clavulanic acid (1000/62.5 mg). *International Journal of Oral and Maxillofacial Surgery* 2007;36(4):321-7.
- Larsen PE: Use of chlorhexidine to prevent alveolar osteitis. *J Oral Maxillofac Surg* 48:1244, 1990
- Leon Arcila ME, Acosta P, Bravo BE, Mena A, Noguera DP, Zúñiga JR. Antibiotic prophylaxis in third molar surgery [Profilaxis antibiologica en cirugía de terceros molares incluidos]. *Revista Estomatología* 2001; Vol. 9, issue 2: 4-13.
- Leung YY, Cheung LK. Coronectomy of the lower third molar is safe within the first 3 years. *J Oral Maxillofac Surg.* 2012;70:1515-22.
- Leung YY, Cheung LK. Long-term morbidities of coronectomy on lower third molar. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2016 Jan;121(1):5-11.
- Leung YY, Cheung LK. Safety of coronectomy versus excision of wisdom teeth: a randomized controlled trial. *Oral Surgery Oral Medicine Oral Pathology Oral Radiology. Endodontics* 2009; Vol. 108, issue 6:821-7.
- Leung YY, McGrath C, Cheung LK. Trigeminal neurosensory deficit and patient reported outcome measures: the effect on quality of life. *PLoS One.* 2013 Oct 29;8(10):e77391.
- Li ZB, Qu HL, Zhou LN, Tian BM, Chen FM. Influence of Non-Impacted Third Molars on Pathologies of Adjacent Second Molars: A Retrospective Study. *J Periodontol.* 2017 May;88(5):450-456.
- Liedholm R, Knutsson K, Lysell L, Rohlin M, Brickley M, Shepherd JP. The outcomes of mandibular third molar removal and non-removal: a study of patients' preferences using a multi-attribute method. *Acta Odontol Scand.* 2000 Dec;58(6):293-8.
- Liedholm R, Knutsson K, Lysell L, Rohlin M, Brickley M, Shepherd J. Third molar treatment outcome: a comparison of patients' preferences in Sweden and Wales. *Br Dent J.* 2005 Sep 10;199(5):287-91.
- Liversidge HM, Marsden PH. Estimating age and the likelihood of having attained 18 years of age using mandibular third molars. *Br Dent J.* 2010 Oct 23;209(8):E13.
- Lodi G, Figini L, Sardella A, Carrassi A, Del Fabbro M, Furness S. Antibiotics to prevent complications following tooth extractions. *Cochrane Database of Systematic Reviews* 2012, Issue 11. Art.No.:CD003811.
- López-Cedrún JL, Pijoan JI, Fernández S, Santamaria J, Hernandez G. Efficacy of amoxicillin treatment in preventing postoperative complications in patients undergoing third molar surgery: a prospective, randomized, double-blind controlled study. *Journal of Oral and Maxillofacial Surgery* 2011;69:e5-e14.
- Ludlow, J.B., et al., Effective dose of dental CBCT-a meta analysis of published data and additional data for nine CBCT units. *Dentomaxillofac Radiol*, 2015. 44(1): p. 20140197.
- Ludlow, J.B., L.E. Davies-Ludlow, and S.C. White, Patient risk related to common dental radiographic examinations: the impact of 2007 International Commission on Radiological Protection recommendations regarding dose calculation. *J Am Dent Assoc*, 2008. 139(9): p. 1237-43.
- Malkawi Z, Al-Omiri MK, Khraisat A. Risk indicators of postoperative complications following surgical extraction of lower third molars. *Med Princ Pract.* 2011;20(4):321-5.
- Marcussen KB, Laulund AS, Jørgensen HL, Pinholt EM. A Systematic Review on Effect of Single-Dose Preoperative Antibiotics at Surgical Osteotomy Extraction of Lower Third Molars. *J Oral Maxillofac Surg.* 2016 Apr;74(4):693-703.
- Monaco G, Tavernese L, Agostini R, Marchetti C. Evaluation of antibiotic prophylaxis in reducing postoperative infection after mandibular third molar extraction in young patients. *Journal of Oral and Maxillofacial Surgery* 2009;67(7):1467-72.
- Matzen LH, Christensen J, Hintze H, Schou S, Wenzel A. Diagnostic accuracy of panoramic radiography, stereo-scanography and cone beam CT for assessment of mandibular third molars before surgery. *Acta Odontol Scand.* 2013 Nov;71(6):1391-8.
- Matzen LH, Hintze H, Spin-Neto R, Wenzel A. Reproducibility of mandibular third molar assessment comparing two cone beam CT units in a matched pairs design. *Dentomaxillofac Radiol.* 2013b;42(10):20130228.
- Matzen LH, Schropp L, Spin-Neto R, Wenzel A. Use of cone beam computed tomography to assess significant imaging findings related to mandibular third molar impaction. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2017 Aug 2. pii: S2212-4403(17)31010-6.
- Matzen LH, Wenzel A. Efficacy of CBCT for assessment of impacted mandibular third molars: a review - based on a hierarchical model of evidence. *Dentomaxillofac Radiol.* 2015;44(1):20140189.
- McArdle LW, Renton TF. Distal cervical caries in the Mandibular second molar: An indication for the prophylactic removal of the third molar? *Br J Oral Maxillofac Surg* 2005;44:42-5.
- Metin M, Tek M, Sener I: Comparison of two chlorhexidine rinse protocols on the incidence of alveolar osteitis following the surgical removal of impacted third molars. *JContempDent Pract* 7:79, 2006
- Mocan A, Kisinisci R, Ucok C. Stereophotogrammetric and clinical evaluation of morbidity after removal of lower third molars by two different surgical techniques. *Journal of Oral and Maxillofacial Surgery* 1996;54(2):171-5.
- Monaco G, de Santis G, Gatto MR, Corinaldesi G, Marchetti C. Coronectomy: a surgical option for impacted third molars in close proximity to the inferior alveolar nerve. *J Am Dent Assoc.* 2012;143:363-9.
- Monaco G, De Santis G, Pulpito G, Gatto MR, Vignudelli E, Marchetti C. What Are the Types and Frequencies of Complications Associated With Mandibular Third Molar Coronectomy? A Follow-Up Study. *J Oral Maxillofac Surg.* 2015 Jul;73(7):1246-53.

- Mukherjee S, Vikraman B, Sankar D, Veerabahu MS. Evaluation of Outcome Following Coronectomy for the Management of Mandibular Third Molars in Close Proximity to Inferior Alveolar Nerve. *J Clin Diagn Res.* 2016 Aug;10(8):ZC57-62.
- Nageshwar N. Comma incision for impacted mandibular third molars. *Journal of Oral and Maxillofacial Surgery* 2002; 60(12):1506-9.
- Naichuan Su et al. Predictive Value of Panoramic Radiography for Injury of Inferior Alveolar Nerve After Mandibular Third Molar Surgery. *J Oral Maxillofac Surg* -:1-17, 2016 (in press).
- Nemcovsky CE, Libfeld H, Zubery Y. Effect of non-erupted 3rd molars on distal roots and supporting structures of approximal teeth. A radiographic survey of 202 cases. *J Clin Periodontol.* 1996 Sep;23(9):810-5.
- Nunn ME, Fish MD, Garcia RI, Kaye EK, Figueroa R, Gohel A, Ito M, Lee HJ, Williams DE, Miyamoto T. Retained asymptomatic third molars and risk for second molar pathology. *J Dent Res.* 2013 Dec;92(12):1095-9. doi: 10.1177/0022034513509281. Epub 2013 Oct 16.
- Oenning AC, Melo SL, Groppo FC, Haiter-Neto F. Mesial inclination of impacted third molars and its propensity to stimulate external root resorption in second molars--a cone-beam computed tomographic evaluation. *J Oral Maxillofac Surg.* 2015 Mar;73(3):379-86.
- Osunde OD, Adebola RA, Saheeb BD. A comparative study of the effect of suture-less and multiple suture techniques on inflammatory complications following third molar surgery. *International Journal of Oral and Maxillofacial Surgery* 2012; Vol. 41, issue 10:1275-9.
- Osunde OD, Saheeb BD, Adebola RA. Comparative study of effect of single and multiple suture techniques on inflammatory complications after third molar surgery. *Journal of Oral and Maxillofacial Surgery* 2011;69:971-6.
- Ozeç I, Hergüner Siso S, Taşdemir U, Ezirganli S, Göktolga G. Prevalence and factors affecting the formation of second molar distal caries in a Turkish population. *Int J Oral Maxillofac Surg.* 2009 Dec;38(12):1279-82.
- Padhye MN, Dabir AV, Girotra CS, Pandhi VH. Pattern of mandibular third molar impaction in the Indian population: a retrospective clinico-radiographic survey. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2013 Sep;116(3):e161-6.
- Pasqualini D, Cocero N, Castella A, Mela L, Bracco P. Primary and secondary closure of the surgical wound after removal of impacted mandibular third molars: a comparative study. *International Journal of Oral and Maxillofacial Surgery* 2005;34(1):52-7.
- Pasupathy S, Alexander M. Antibiotic prophylaxis in third molar surgery. *Journal of Craniofacial Surgery* 2011;22: 551-3.
- Petersen LB, Olsen KR, Christensen J, Wenzel A. Image and surgery-related costs comparing cone beam CT and panoramic imaging before removal of impacted mandibular third molars. *Dentomaxillofac Radiol.* 2014;43(6):20140001.
- Petersen LB, Vaeth M, Wenzel A. Neurosensoric disturbances after surgical removal of the mandibular third molar based on either panoramic imaging or cone beam CT scanning: A randomized controlled trial (RCT). *Dentomaxillofac Radiol.* 2016;45(2):20150224.
- Pogrel MA, Lee JS, Muff DF. Coronectomy: a technique to protect the inferior alveolar nerve. *J Oral Maxillofac Surg.* 2004;62:1447-52.
- Polat HB, Ozan F, Kara I, Ozdemir H, Ay S. Prevalence of commonly found pathoses associated with mandibular impacted third molars based on panoramic radiographs in Turkish population. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008 Jun;105(6):e41-7.
- Praveen G, Rajesh P, Neelakandan RS, Nandagopal CM. Comparison of morbidity following the removal of mandibular third molar by lingual split, surgical bur and simplified split bone technique. *Indian Journal of Dental Research* 2007;18(1):15-8.
- Ragno JR Jr, Szkutnik AJ: Evaluation of 0.12% chlorhexidine rinse on the prevention of alveolar osteitis. *Oral Surg Oral Med Oral Pathol* 72:524, 1991
- Rakprasitkul S, Pairuchvej V. Mandibular third molar surgery with primary closure with a drain. *International Journal of Oral and Maxillofacial Surgery* 1997;26(3): 187-90.
- Refo'a Y, Ouatik N, Golchin F, Mahboobi N. Comparing primary and secondary wound healing discomfort after mandibular third molar surgery: a randomized, double blind clinical trial. *General Dentistry* 2011; Vol. 59, issue 4:310-3.
- Renton T, Hankins M, Sproate C, McGurk M. A randomised controlled clinical trial to compare the incidence of injury to the inferior alveolar nerve as a result of coronectomy and removal of mandibular third molars. *British Journal of Oral and Maxillofacial Surgery* 2005;43(1): 7-12.
- Ritzau M, Hillerup S, Branbjerg PE, Ersbol BK. Does metronidazole prevent alveolitis sicca dolorosa? A doubleblind, placebo-controlled clinical study. *International Journal of Oral and Maxillofacial Surgery* 1992;21(5): 299-302.
- Rodríguez Sánchez F, Rodríguez Andrés C, Arteagoitia Calvo I. Does Chlorhexidine Prevent Alveolar Osteitis After Third Molar Extractions? Systematic Review and Meta-Analysis. *J Oral Maxillofac Surg.* 2017 Jan 13. pii:S0278-2391(17)30020-4. doi: 10.1016/j.joms.2017.01.002
- Rodríguez-Perez M, Bravo-Perez M, Sanchez-Lopez JD, et al: Effectiveness of 1% versus 0.2% chlorhexidine gels in reducing alveolar osteitis from mandibular third molar surgery: A randomized, double-blind clinical trial. *Med Oral Patol Oral Cir Bucal* 18: e693, 2013
- Roode GJ, Butow K. An alternative surgical flap design for impacted third molars: a comparison of two different surgical techniques. *SADJ* 2010;65(6):246, 248-51.
- Rubio-Palau J, Garcia-Linares J, Hueto-Madrid JA, et al: Effect of intra-alveolar placement of 0.2% chlorhexidine bioadhesive gel on the incidence of alveolar osteitis following the extraction of mandibular third molars. A double-blind randomized clinical trial. *Med Oral Patol Oral Cir Bucal* 20:e117, 2015
- Rullo R, Addabbo F, Papaccio G, D'Aquino R, Festa VM. Piezoelectric device vs. conventional rotative instruments in impacted third molar surgery: relationships between surgical difficulty and postoperative pain with histological evaluations. *Journal of Cranio-Maxillo-Facial Surgery* 2013; 41(2):e33-8. [PUBMED: 22890087]

- Saglam AA. Effects of tube drain with primary closure techniques on postoperative trismus and swelling after removal of fully impacted mandibular third molars. *Quintessence International* 2003;34(2):143-7.
- Sandhu A, Sandhu S, Kaur T. Comparison of two different flap designs in the surgical removal of bilateral impacted mandibular third molars. *International Journal of Oral and Maxillofacial Surgery* 2010;39(11):1091-6.
- Sekhar CH, Narayanan V, Baig MF. Role of antimicrobials in third molar surgery: prospective, double blind, randomized, placebo-controlled clinical study. *British Journal of Oral & Maxillofacial Surgery* 2001; Vol. 39, issue 2:134-7. Absi EG, Shepherd JP. A comparison of morbidity following the removal of lower third molars by lingual split and surgical bur methods. *International Journal of Oral and Maxillofacial Surgery* 1993;22(3):149-53.
- Shin SM, Choi EJ, Moon SY. Prevalence of pathologies related to impacted mandibular third molars. *Springerplus*. 2016 Jun 29;5(1):915.
- Shiratori K, Nakamori K, Ueda M, Sonoda T, Dehari H. Assessment of the shape of the inferior alveolar canal as a marker for increased risk of injury to the inferior alveolar nerve at third molar surgery: a prospective study. *J Oral Maxillofac Surg*. 2013 Dec;71(12):2012-9.
- Shugars DA, Elter JR, Jacks MT, White RP, Phillips C, Haug RH, Blakey GH. Incidence of occlusal dental caries in asymptomatic third molars. *J Oral Maxillofac Surg*. 2005 Mar;63(3):341-6.
- Srinivas DR. Effect of Surgical Tube Drain with Primary Closure Technique after Removal of Impacted Mandibular Third Molars - a Clinical Study (Dissertation). Karnataka, Bangalore: Rajiv Gandhi University of Health Sciences. Karnataka, Bangalore, India: Rajiv Gandhi University of Health Sciences, 2006:1-51.
- Srivastava N, Shetty A, Goswami RD, Apparaju V, Bagga V, Kale S. Incidence of distal caries in mandibular second molars due to impacted third molars: Nonintervention strategy of asymptomatic third molars causes harm? A retrospective study. *Int J Appl Basic Med Res*. 2017 Jan-Mar;7(1):15-19.
- Stathopoulos P, Mezitis M, Kappatos C, Titsinides S, Stylogianni E. Cysts and tumors associated with impacted third molars: is prophylactic removal justified? *J Oral Maxillofac Surg*. 2011 Feb;69(2):405-8.
- Sweet JB, Butler MS, Drager JL. Effects of lavage techniques with third molar surgery. *Oral Surgery, Oral Medicine, and Oral Pathology* 1976;41(2):152-68.
- Tantanapornkul W, Okouchi K, Fujiwara Y, Yamashiro M, Maruoka Y, Ohbayashi N, Kurabayashi T. A comparative study of cone-beam computed tomography and conventional panoramic radiography in assessing the topographic relationship between the mandibular canal and impacted third molars. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007 Feb;103(2):253-9.
- Tjernberg A: Influence of oral hygiene measures on the development of alveolitis sicca dolorosa after surgical removal of mandibular third molars. *Int J Oral Surg* 8:430, 1979
- Toedtling V, Coulthard P, Thackray G. Distal caries of the second molar in the presence of a mandibular third molar - a prevention protocol. *Br Dent J*. 2016 Sep 23;221(6):297-302.
- Torres-Lagares D, Gutierrez-Perez JL, Infante-Cossio P, et al: Randomized, double-blind study on effectiveness of intra-alveolar chlorhexidine gel in reducing the incidence of alveolar osteitis in mandibular third molar surgery. *Int J Oral Maxillofac Surg* 35:348, 2006
- Ventä I, Kylätie E, Hiltunen K. Pathology related to third molars in the elderly persons. *Clin Oral Investig*. 2015 Nov;19(8):1785-9.
- Ventä I, Meurman JH, Murtomaa H, Turtola L. Effect of erupting third molars on dental caries and gingival health in Finnish students. *Caries Res*. 1993;27(5):438-43.
- Ventä I, Vehkalahti MM, Huuonen S, Suominen AL. Signs of disease occur in the majority of third molars in an adult population. *Int J Oral Maxillofac Surg*. 2017 Aug 16. pii: S0901-5027(17)31533-3.
- von Wowern N, Nielsen HO. The fate of impacted lower third molars after the age of 20. A four-year clinical follow-up. *Int J Oral Maxillofac Surg*. 1989 Oct;18(5):277-80.
- Wang D, He X, Wang Y, Li Z, Zhu Y, Sun C, Ye J, Jiang H, Cheng J. External root resorption of the second molar associated with mesially and horizontally impacted mandibular third molar: evidence from cone beam computed tomography. *Clin Oral Investig*. 2017 May;21(4):1335-1342.
- Welch VA, Akl EA, Guyatt G, Pottie K, Eslava-Schmalbach J, Ansari MT, de Beer H, Briel M, Dans T, Dans I, Hultcrantz M, Jull J, Katikireddi SV, Meerpohl J, Morton R, Mosdol A, Petkovic J, Schünemann HJ, Sharaf RN, Singh JA, Stanev R, Tonia T, Tristan M, Vitols S, Watine J, Tugwell P. GRADE equity guidelines 1: considering health equity in GRADE guideline development: introduction and rationale. *J Clin Epidemiol*. 2017 Oct;90:59-67.
- Wijs RW, Karssemakers LH, Becking AG. [Coronectomy. An alternative for complete removal of the third molar in the lower jaw]. *Ned Tijdschr Tandheelkd*. 2010 Jun;117(6):337-40.
- Winter G.B. Impacted mandibular third molars. St Louis: American Medical Book Co.; 1926. p. 241-79.
- Wu TH, Lin WC, Chen WK, Chang YC, Hwang JJ. Predicting cancer risks from dental computed tomography. *J Dent Res*. 2015 Jan;94(1):27-35.
- Xavier RL, Vasconcelos BC, Caubi AF, Porto GG, Maurette MA. Passive drainage through the vestibular oblique incision in impacted inferior third molar surgery: a preliminary study. *Acta Odontologica Latinoamericana* 2008; Vol. 21, issue 1:57-63.
- Yamalik K, Bozkaya S. The predictivity of mandibular third molar position as a risk indicator for pericoronitis. *Clin Oral Investig*. 2008 Mar;12(1):9-14.
- Yamaoka M, Furusawa K, Ikeda M, Hasegawa T. Root resorption of mandibular second molar teeth associated with the presence of the third molars. *Aust Dent J*. 1999 Jun;44(2):112-6.
- Zandi M, Shokri A, Heidari A, Masoud Peykar E. Objectivity and reliability of panoramic radiographic signs of intimate relationship between impacted mandibular third molar and inferior alveolar nerve. *Oral Maxillofac Surg*. 2015 Mar;19(1):43-8.

Zhou J, Hu B, Liu Y, Yang Z, Song J. The efficacy of intra-alveolar 0.2% chlorhexidine gel on alveolar osteitis: a meta-analysis. Oral Dis. 2016 Aug 1. doi: 10.1111/odi.12553